

****FULL TITLE****

*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***

****NAMES OF EDITORS****

Infrared Properties of Cataclysmic Variables in the 2MASS All Sky Data Release

D. W. Hoard¹, C. S. Brinkworth^{1,2}, S. Wachter¹

¹*Spitzer Science Center, California Institute of Technology, Pasadena, CA 91125, USA*

²*Department of Physics and Astronomy, University of Southampton, Highfield, Southampton SO17 1BJ, UK*

Abstract. We present results from our analysis of the near-infrared (J , H , and K_s) photometry for all cataclysmic variables (CVs) from the catalog of Downes et al. (2001) that are detected in the (final) All Sky Data Release (ASDR) from the Two Micron All Sky Survey (2MASS).

1. Introduction

CVs have been “traditionally” observed primarily at short wavelengths because accretion-generated luminosity, which peaks in the optical-ultraviolet, dominates the radiated energy of most systems. Hence, relatively little is known about their infrared (IR) properties. Investigating CVs in the IR contributes to the understanding of system components that are expected to radiate at these wavelengths, such as the cool outer disk, accretion stream, and secondary star. In Hoard et al. (2002), we presented an initial study of the group IR properties of CVs using the 2MASS 2nd Incremental Data Release (2IDR), drawn from 525 valid targets located in the 2IDR sky coverage. We present here the preliminary results obtained from extending this study to the 2MASS ASDR.

2. Data Properties

Our input target list for the 2MASS ASDR consisted of the 1320 valid CVs listed in Downes et al. (2001) as of 02 February 2004. Each CV was securely identified using optical finding charts from Downes et al. (2001) and/or other literature sources before matching it to an object in the 2MASS ASDR images and point source catalog. We could not recover the CV for 239 targets (typically CVs in crowded star clusters, old novae near the Galactic plane, etc.) and could not securely identify the IR counterpart for 104 (typically in regions where the IR field is very crowded). The locations of another 376 CVs were securely identified, but these systems were too faint to be detected by 2MASS. The remaining 601 CVs were detected by 2MASS; of these, we classify 362 as “good” detections (1σ uncertainties less than 0.1 mag for J , H , and K_s), 165 as “moderate” detections (1σ uncertainty larger than 0.1 mag for at least one of J , H , or K_s), and 74 as “poor” detections (lacking a formal uncertainty for at least one of J , H , or K_s , indicating that the target is near the 2MASS faint detection limit).




Figure 1. (left) IR color-color diagram for all CVs classified as “good” detections (see text) in 2MASS. (right) IR color-magnitude diagram for CVs detected in 2MASS, with distances determined from trigonometric parallax.

3. Results

Figure 1 (left) shows the IR color-color diagram of all “good” CV detections from 2MASS. The loci of the main sequence (MS), L dwarfs, and giant stars are shown as cross-hatched regions labelled with spectral types (offset horizontally) in normal (MS, L dwarfs) and italic (giants) fonts. The CVs located in outlying color regions (e.g., L dwarf region, lower right quadrant, etc.) may be mis-identified non-CVs or potentially interesting systems. Figure 1 (right) shows the IR color-magnitude diagram for CVs with distances determined from trigonometric parallax (e.g., Harrison et al. 1999; Thorstensen 2003; etc.). The object closest to the L dwarf region is EF Eri, a CV that is believed to contain a brown-dwarf-like secondary star (Howell & Ciardi 2001). We are currently updating the input target sample to include all CVs in Downes et al. (2001) up to 01 September 2004 for final analysis of the IR properties of CVs, to appear in a future publication.

Acknowledgments. This research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, and is based on work supported by the National Aeronautics and Space Administration under an ADP grant issued through the Office of Space Science. CSB acknowledges support from the Spitzer Science Center Visiting Graduate Student Program. This work uses data products from 2MASS, a joint project of the U. of Massachusetts and IPAC/Caltech, funded by NASA and the NSF.

References

- Downes, R. A., et al. 2001, *PASP*, 113, 764
- Harrison, T. E., et al. 1999, *ApJ*, 515, L93
- Hoard, D. W., Wachter, S., Clark, L. L., & Bowers, T. P. 2002, *ApJ*, 565, 511
- Howell, S. B. & Ciardi, D. R. 2001, *ApJ*, 550, L57
- Thorstensen, J. R. 2003, *AJ*, 126, 3017